

Journal of Spices and Aromatic Crops
Vol. 20 (1) : 01–08 (2011)


Indian Society for Spices

Use of indigenous plant products for management of pests and diseases of spices and condiments: Indian perspective

R T Gahukar

*Arag Biotech Private Limited
Plot 220, Reshimbag, Nagpur-440 009, India.
E-mail: rtgahukar@gmail.com*

Received 24 July 2010; Revised 18 December 2010; Accepted 10 January 2011

Abstract

Plant derived preparations and formulated commercial products have great potential for eco-friendly management of pests and diseases of spice and condiment crops in India. These products act as antifeedant, growth regulator, repellent and a direct mortality factor against pests and as inhibitor of sporulation and development of hyphae of fungal pathogens. However, further research on field application, residual toxicity, synergistic action and hazardous effects on non-target organisms are needed.

Keywords: condiments, diseases, IPM, pests, plant products, spices.

Introduction

The productivity of most of the spice and condiment crops is considerably low in India due to various factors among which infestation by pests and pathogens is a major factor. Several species of insects, mites, nematodes and pathogens attack these crops in the field and also during storage. Synthetic pesticides have been used extensively earlier to reduce crop damage and increase crop profitability leading to pesticide residues in the produce, harmful side effects on beneficial organisms, mortality of pollinators, environmental pollution and resistance to pesticides. Since spices are high-value and export-oriented commodities, levels of pesticide residues are to be kept well below tolerance limits in view of the stringent standards set by the importing countries (Devasahayam 2005). Thus, the growing

concern over pesticide misuse (excessive and indiscriminate) has led to the development of IPM schedules for sustainable management of pests and diseases in recent years. Plant products form an important component in the development of IPM schedules.

Among indigenous plant species, neem (*Azadirachta indica* A. Juss.) has been extensively studied for bioactive constituents in different plant parts and over 50 commercial products containing the major bioactive isomer, Azadirachtin (AZ) are now available in the market for general use (Gahukar 1998; Shanker & Parmar 1999). Farmers use local traditional preparations such as, neem seed kernel extract (NSKE), neem seed extract (NSE), neem leaf extract (NLE), neem crude extract (NCE), neem cake (NC) and neem oil (NO) for the management of insect pests. Different modes of action

(antifeedant, repellent, insect growth regulator, oviposition deterrent and pesticidal) of the limonoids, terpenoids, flavonoids and other alkaloids present in plant products have been known through research findings (Gahukar 1995). Apart from neem, *karanj/pongam* (*Pongamia pinnata* (L.) Pierre) and *nirgudi* (*Vitex negundo* L.) are also being used in traditional preparations (crude extract, water extract, cake) and commercial products. Therefore, it is now possible to minimize the application of hazardous chemical pesticides. This paper reviews the information available on plant products used on spice and condiment crops in India and outlines suggestions for further improvement in this aspect.

Betelvine (*Piper betle* L.)

Sap sucking insects (aphids, thrips, whiteflies and scale insects) and mites are major pests and leaf spots, blight and rots are common diseases. In some places, infestations of nematodes have also been reported. Among the several species of mites infesting betelvine, the red mite (*Tetranychus neocaledonicus* Andre) is a major pest. The nymphs and adult mites suck sap from young leaves that turn yellow and drop. Nakat *et al.* (2000) tested the bioefficacy of NSKE (5%), tobacco decoction (2%) and dicofol 18.5 EC (0.05%) under laboratory and field conditions. All treatments provided satisfactory control but plant products were recommended as they are safer to predators. A spray of NSKE (5%) or NO (2%) has been recommended against linear scale (*Lepidosaphes cornutus* Ramkrishnan) (Chandra & Sagar 2004) on betelvine.

Neem leaf extract (2%) or extract of NC (0.5%) was recommended against *Phytophthora capsici* Leonian causing foot rot disease. Mixing of soil with chopped leaves of *Calotropis* sp. and neem (Chandra & Sagar 2004), or a mixture of neem cake and saw dust (Acharya & Padhi 1988) at planting time has been suggested against nematodes, *Meloidogyne incognita* (Kofoid & White) and *Hoplolaimus indicus* Sher. Neem cake applied @ 2 t ha⁻¹ also controlled wilt disease caused by a fungal complex (Lal 1993).

Black pepper (*Piper nigrum* L.)

The major pests of black pepper include *pollu* beetle, scale insects, and nematodes and major diseases include *Phytophthora* foot rot and slow wilt. The *pollu* beetle (*Longitarsus nigripennis* Mots.) is the most destructive insect pest of black pepper. The beetles feed on leaves, growing shoot tips and tender spikes. The grubs bore into growing shoot tips, spikes and berries and feed on internal contents. Bioassay of plant based pesticides indicated that commercial neem products such as Repelin®, Neemgold® and Neemazal-F® (2%-3%), and leaf extracts (3%) of *Strychnos nux-vomica* L., *Chromolaena odorata* L. and *Annona squamosa* L. caused 90% antifeeding effect in laboratory bioassays and Neemgold® was the most effective treatment in the field (Devasahayam & Anandaraj 1997; Devasahayam & Leela 1997). Root bark extract of *Uvaria narum* (Dunal) Blume and *U. hookeri* King also caused >90% feeding inhibition at 0.01%-0.05% concentrations in the laboratory (Babu *et al.* 1996). For effective pest management, Devasahayam & Koya (1999) recommended regulation of shade during May-June, spraying with quinalphos 25 EC (0.05%) during July followed by Neemgold® (0.6%) during August-October at 21-day intervals. In case of scale insects (*Lepidosaphes piperis* Green and *Aspidiotus destructor* Sign.), spot application of monocrotophos (0.1%) or dimethoate (0.1%) followed by Neemgold® (0.6%) proved to be effective (Selvan *et al.* 1996; Devasahayam & Koya 1999).

Extracts of five plant species on growth stages of *P. capsici* were evaluated in laboratory bioassays among which *C. odorata* caused 100% inhibition of mycelial growth, sporangial production, zoospore production, release and germination (Anandaraj & Leela 1996). Essential oil of allspice (*Pimenta dioica* L. Merr.) leaves and its major constituent eugenol exhibited promising nematocidal activity above 660 µg ml⁻¹ against root knot nematode *M. incognita* (Leela & Ramana 2000).

Cardamom (*Elettaria cardamomum* Maton)

Thrips, shoot and capsule borer, root grubs and hairy caterpillars are considered as major pests; capsule rot (*azhukal* disease), leaf blight, necrosis and mosaic (*katte* disease) are major diseases.

Cardamom plants are heavily infested by thrips (*Sciothrips cardamomi* Ramk.), resulting in shedding of flowers and immature capsules and formation of malformed capsules with corky, scab-like encrustation. Since plant products such as, Neemgold® (3%), Econeem® (0.03%), NSKE (4%) and NO (3%) could not control the pest on capsules, many chemical pesticides have generally been recommended (Naik *et al.* 2004). In nursery, young larvae of shoot and capsule borer (*Conogethes punctiferalis* Guen.) bore into and feed on the internal contents of unopened leaf buds, panicles and immature capsules while later instars bore into pseudostems and cause dead hearts due to drying of growing shoot. Amongst the neem products experimented in the field, Neemgold® (3%) and NO (3%) were promising (Naik *et al.* 2004). Colonies of aphid (*Pentalonia nigronervosa* f. *caladii* Van der Goot) remain inside leaf sheaths and suck plant sap in the nursery and plantation. These species are also a vector of the virus that causes the dreaded *katte* disease. Laboratory bioassays and field trials revealed that seed extract of *A. squamosa*, leaf extract of *Lawsonia inermis* L. and *Leucas aspera* (Willd.) Spreng., rhizome extract of *Acorus calamus* L. and two neem products (Bioneem®, Margolin®) significantly affected the settling and colonization behaviour and multiplication of aphids in cardamom (Venugopal 1999). Gopakumar *et al.* (1996) recommended NO (3%) or a water extract (5%) of *Lantana camara* L. leaves against whiteflies, *Dialeurodes cardamomi* David & Subr. which cause depletion of sap from leaves. These products act as repellents and prevent their population build-up in the plantation.

Chilli (*Capsicum annuum* L.)

The chilli crop is mainly infested by thrips, aphids, whiteflies, jassids and mealybugs on

foliage and pod borer on fruits. Attack of root grubs and yellow mite is also severe in certain seasons. Anthracnose, damping off, bacterial wilt, root rot, grey mould, bacterial brown rot and blossom end rot are common diseases; viral diseases include leaf curl, mosaic and yellow dwarf.

The sucking pests suck sap from young and mature leaves and shoots resulting in curling of leaves, wilting and yellowing of young shoots, drying of flowers, and stunted growth of seedlings. Whiteflies also act as vectors of viral diseases such as, chilli mottle virus and tomato leaf curl. Sprays of neem-based commercial products such as, Achook®, Nimin®, Nimarin® (1%) applied @ 62.5 g a.i. ha⁻¹ after mixing with urea @ 62 kg ha⁻¹ resulted in less mortality of whitefly, *Bemisia tabaci* Genn. than with monocrotophos 36 SL (0.05%) (Saha *et al.* 2005). Therefore, a mixture of monocrotophos and Achook® or Nimin® and urea (Keisa & Varatharajan 1995) or monocrotophos (0.03%) alone had been suggested by Saha *et al.* (2005). The spraying against whitefly could also prevent the spread of leaf curl virus (Saha *et al.* 2005). Against broad mite (*Polyphagotarsonemus latus* (Banks)), sprays of Groneem® (0.015 EC) or NSKE (5%) were not effective, whereas a spray of synthetic miticide, Omite® 300W-WP (0.228%) resulted in 74% pest mortality (Kumar *et al.* 2005 b). Chilli thrips (*Scirtothrips dorsalis* Hood), is a major pest throughout the year. Although neem products controlled the pest effectively (Keisa & Varatharajan 1995), sprays of imidacloprid 200 SL (0.5 ml⁻¹) mixed with NO or *P. pinnata* oil (0.2%) were more effective (IIHR 2008). Two major pod borers namely, *Helicoverpa armigera* (Hbn.) and *Spodoptera litura* (F.) destroy fruit buds and bore into developing fruits and devour the inner content making the fruit unsuitable for marketing. An alkaloid derived from *Saphora* spp. (sold as Oxymatrine® 98%) reduced fruit damage and was significantly better than synthetic pesticides (endosulfan, chlorpyrifos, carbaryl) (Sekhar & Ram 2004). The gall midge (*Aspondylla capsicici* Barnes) feeds on interior content of flower

bud resulting in the formation of galls and fruit formation is inhibited. By spraying NO (1%) or NCE (5%) mixed with a sticker, Sandovit® (0.05%), during flowering stage, the pest infestation was reduced up to 57% but these treatments were not at par compared to phorate (10%) or carbofuran (3%) granules incorporated into soil @ 1 kg a.i. ha⁻¹ (David *et al.* 1990). In order to reduce the application cost of treatment against sucking pests and pod borers, Rao *et al.* (1998; 1999 a, b) suggested the following IPM module: (i) incorporation of NC into soil before transplanting @ 500 kg ha⁻¹ (ii) dipping seedling roots in NO (1%) (iii) weekly sprays of NO (1%); these treatments were at par with dimethoate (0.03%) in bioefficacy. Jain & Ameta (2006) suggested two sprays of imadacloprid 200 SL (750 ml ha⁻¹) followed by betacyfluthrin 0.25 EC (750 ml ha⁻¹) because Nimbecidine® (0.3%) was only moderately effective. There was increase in yield and cost: benefit ratio due to significant reduction in pest population and plant damage in all treatments except control.

Pest injury to fruits and larval frass/excreta left inside, attracts *Aspergillus flavus* (Link.) Fres. which is responsible for fruit rotting. Therefore, efforts were made to inhibit development of fungal hyphae by spraying NSKE, Nimbecidine® or oil of *P. pinnata* (5%) in the field (Kumar *et al.* 2005 a). In the field, the disease incidence was reduced from 98% to 24%-30% with sprays of water extract (5%) of *Allium sativum* L., *Datura metel* L., *Eucalyptus globules* Labill. or *Prosopis juliflora* (Sw.) DC, though carbendazim 50 WP (0.1%) spray was the most effective treatment (18% incidence) (Raj *et al.* 2006). Root rot caused by *Alternaria tenuis* Nees was checked by spraying (100-115 days after sowing) with water extract (10%) of *Aegle marmelos* (L.) Corr. or *Prosopis juliflora* (Sw.) DC (Sivaprakasam & Seetharaman 1997). Choudhary *et al.* (2000) experimented soil incorporation of a mixture of *Trichoderma viride* (Pers.) (15 g kg⁻¹ soil) + chestnut compound (25 ml kg⁻¹ soil) + NC (100 g kg⁻¹ soil) and obtained 97% control of wilt disease caused

by *Sclerotium rolfsii* Sacc. and cent per cent survival of the seedlings.

Several neem-based preparations have been tested successfully against root nematodes, especially, those belonging to the genera *Meloidogyne*, *Heterodera* and *Globodera*. Water extract of neem leaves or neem cake for root dipping; NO for seed coating and root dipping, and commercial products for root dipping were promising against these nematode species (Akhtar 2000). Recently, Rao (2007) recommended soil application of NC or *P. pinnata* @ 500 g m² in the nursery a fortnight before sowing, and @ 500 kg ha⁻¹ in the field.

Cinnamon (*Cinnamomum zeylanicum* Blume)

Fruit borer on fruits and leaf miner on foliage are major pests. Major diseases of cinnamon are brown root rot, grey leaf spot, stripe canker, leaf blight and pink disease. In the laboratory, the growth and development *Phytophthora cinnamomi* Rands causing root rot could successfully be inhibited by 5% water extract of eucalyptus (*Eucalyptus* sp.) leaves (Sivasithambaram *et al.* 1981).

Cumin (*Cuminum cyminum* L.)

Aphids are serious pests of cumin. Fungal diseases such as leaf blight, wilt, powdery mildew and rots are commonly observed. Nymphs and adult aphids (*Myzus persicae* Sulz.) suck sap from young and mature leaves which subsequently curl, dry and shed off. NO (4%) was effective in achieving cent per cent control and reducing plant injury (Vir & Yadav 2007).

Curry leaf (*Murraya koenigii* (L.) Spreng.)

Psyllid and lemon butterfly are major pests; *Diaphorina* pink disease and leaf spot are major diseases of curry leaf. Larvae of lemon butterfly (*Papilio demoleus* (L.)) defoliate the plants throughout the year. When Nimbecidine® (0.3% AZ) was sprayed on grown up trees, the neem product acted as antifeedant, insect growth regulator and pesticide against all larval instars (Ramarethinam & Loganathan 2001).

Ginger (*Zingiber officinale* Rosc.) & Turmeric (*Curcuma longa* L.)

Shoot borer, leaf roller, thrips and rhizome scale are major insect pests in the field. The cigarette beetle and drug store beetle are major pests on stored rhizomes. Nematode infestation occurs occasionally. Among the diseases, collar rot, soft/rhizome rot, bacterial wilt, leaf spot and leaf blotch are major diseases.

The shoot borer (*Conogethes punctiferalis* Guen.) is the most destructive pest of ginger and turmeric. The larvae bore into pseudostems and feed on central growing shoot causing dead hearts. Evaluation of NO (1%) and Nimbecidine® (1%) indicated that these products were effective only on ginger when sprayed at fortnightly intervals during July-October (IISR 2003). An integrated approach consisting of mulching of neem leaves @ 10 t ha⁻¹ and spraying with NSKE (5%) during high pest incidence resulted in an increase in yield by 50%-72%; and this treatment was as effective as quinalphos (0.05%) in controlling the pest (Lalnuntluanga & Singh 2008). The injury of scale insect (*Aspidiella hartii* Sign.) to rhizomes is seen as encrustations on the rhizomes and severely infested rhizomes wither and dry. Storage of ginger rhizomes in dry leaves of *S. nux-vomica*, *Clerodendron infortunatum* Linn. or *Glycosmis cochinchinensis* Pierre was promising in checking the pest infestation (Devasahayam & Koya 1999; IISR 2003).

In water logged ginger fields, soft rot is a common fungal disease caused by a complex of *Fusarium* spp. and *Pythium* spp. In the laboratory, a water extract (5%) of neem, *Agave americana* L., *Cassia fistula* L., *V. negundo* and *Eucalyptus* sp. reduced radial growth of hyphae of *Fusarium* and *Pythium* (Panday *et al.* 1992; Sharma 1998) whilst in the field, the disease incidence was lowered by about 50% by mulching with neem leaves (Das 1999; Mezhatu *et al.* 2008). The same fungal rot in turmeric could be managed by soil incorporation of NC @ 2 t ha⁻¹ or organic

manure @ 50 kg ha⁻¹ mixed with *T. viride* powder (1 kg ha⁻¹). Both the treatments increased rhizome yield significantly (Ramarethinam & Rajagopal 1999).

Onion (*Allium cepa* L.)

Among the major pests, thrips, mites, cutworms, onion maggot and head borer are most common. Purple blotch, stem *Phyllum* blight, *Colletotrichum* blight, damping off, downy mildew, bacterial brown rot and yellow dwarf disease are common diseases. Leaf blight caused by *Alternaria* spp. becomes serious during later stages of plant growth. In laboratory and field trials, a commercial product containing palmrosa (*Cymbopogon martini* (Roxb.) Wats) oil applied at 0.1% concentration, inhibited hyphal growth and proved as effective as mancozeb (0.25%) (Karthikeyan *et al.* 2006).

Challenges and perspectives

Though plant products have been evaluated against a many insect and nematode pests of spice and condiment crops they are yet to be experimented for the management of many major pests such as mirid bugs (*Disphinctus politus* (Westw.), *D. measarum* Link.) on betelvine; defoliators (*Lnoudera vittata* (Wlk.), *Eupterote* spp., *Arcilasisa plagiata* Mots., *Euproctis lutifacia* (Hamps.) on cardamom and leaf webber (*Udaspes folus* Cram.) on chilli. Among diseases, plant products have been evaluated against a few diseases only.

Water extracts of plants are easily washed off plants due to heavy rains. Also, because of slow effect, poor contact action and less residual toxicity, more frequent applications of plant products than that of chemicals are needed (Gahukar 1995; Lakshmisubramanian *et al.* 1998). Hence farmers must be convinced of extended pest mortality and ecological and economical benefits that they would derive from IPM including use of plant products. When plant products are mixed with each other or other pesticides, the synergism may show better effect than individual treatment. There is also an urgent need to work on cheaper and locally available stickers and

additives to prolong storage period of preparations, increase residual toxicity, improve effectiveness and minimize cost of treatment.

Plant products were found safer to coccinellid beetles, *Chilocorus cicutatus* Gyllen and *C. nigra* (F.) predating on scale insects on black pepper (Devasahayam & Koya 1999), and thrips (*Scolothrips* spp.) and mites (*Amblyseius congoensis* Fain) predating on red mite on betel vine (Nakat *et al.* 2000). Similarly, feeding activity of another coccinellid beetle *Cryptolaemus montrouzieri* Mulsant which is a major predator of *Trialeurodes vaporariorum* (Westw.), was not affected at 1000 ppm of (Simmonds *et al.* 2000). However, in laboratory experiments, growth of two important beneficial fungi, *Metarrhizium anisopliae* (Metch.) Sorokin and *Beauveria bassiana* (Bals.) Vuill., was adversely affected by the toxic effects of plant products such as AZ at 2000 ppm (Gupta *et al.* 2002); NSKE (5%) and water extract (5%) of *Ocimum sanctum* L., *A. sativum*, *A. calamus* and *Tribulus terrestris* L. (Devaprasad *et al.* 1989). Development period of chrysopid larvae was prolonged due to NSKE (5%) and Neemark® containing 1500 ppm of AZ (5 ml L⁻¹) (More *et al.* 2005). Direct toxic effects on other natural enemies, pollinators, honey bees and entomopathogens are required to be studied intensively and only safe indigenous products should be recommended.

The content of biologically active constituents in plants differs as per ecotypes (Senguttuvan *et al.* 2005), genetic diversity (Vir 2007) and climatic conditions (Gupta *et al.* 2010). It is therefore necessary to identify those ecotypes with higher content of AZ and other limonoids. These synthesized commercial products and traditional preparations from such geographical areas must be patented (Gahukar 2003). There is also an industrial perspective for plant products due to abundance of indigenous resources for plant-based pesticides (Suresh *et al.* 2001). Education of farmers on proper methods of collection, storage and preparation of plant based preparations would also help to

popularize plant products for the management of pests and diseases of spice and condiment crops.

References

- Acharya A & Padhi N N 1988 Effect of neem oil cake and saw dust against root-knot nematode, *Meloidogyne incognita* on betelvine (*Piper betle* L.). Indian J. Nematol. 18: 105-206.
- Akhtar M 2000 Nematicidal potential of the neem tree, *Azadirachta indica* (A. Juss). Integr. Pest Manage. Rev. 6: 57-66.
- Anandaraj M & Leela N K 1996 Toxic effect of some plant extracts on *Phytophthora capsici*, the foot rot pathogen of black pepper. Indian Phytopathol. 49: 181-184.
- Babu P B S, Padmaja V & Hishan A 1996 Isodesacetylurarin: Insect antifeedant against *Longitarsus nigripennis* Mots. Indian J. Exp. Biol. 34: 377-379.
- Chandra G & Sagar R L 2004 Harvesting green gold: Cultivation of betelvine in Sundarban. Indian Farm. Dig. 37 (3): 5-7.
- Choudhary A K, Rajaram Reddy D, Chandrasekhara Rao K, Bhupal Reddy T & Prabhakar Reddy I 2000 Integrated management of sclerotial wilt disease of bell pepper (*Capsicum annuum* L.). Indian J. Plant Prot. 28: 15-18.
- Das N 1999 Effect of organic mulching on root knot nematode population, rhizome rot incidence and yield of ginger. Ann. Plant Prot. Sci. 77 (1): 12-14.
- David P M M, Jaysekhar M & Natarajan S 1990 Evaluation of insecticides and botanicals for the control of flower gall midge on chillies. Madras Agric. J. 77: 249-252.
- Devaprasad V, Jayaraj S & Rabindra R J 1989 Effect of certain botanicals on the conidial germination in *Beauveria bassiana* (Bals.) Vuill. J. Biol. Contr. 3: 133.
- Devasahayam S 2005 Potential use of plant products for the management of insect pests of spices. Pestology 29 (11): 14-16.
- Devasahayam S & Anandraj M 1997 Black pepper (*Piper nigrum* L.) research. In: Narwal S S, Pauro P & Bisisa S S (Eds.) Neem in Sustainable Agriculture (pp. 117-122). Scientific Publishers, Jodhpur.
- Devasahayam S & Koya K M A 1999 Integrated management of insect pests of spices.

- Indian J. Arecanut Spices Med. Plants 1: 19-23.
- Devasahayam S & Leela N K 1997 Evaluation plant products for antifeedant activity against pollu beetle (*Longitarsus nigripennis* Motschutsky), a major pest of black pepper. In: Reddy P P, Kumar N K & Verghese A (Eds.) Advances in IPM for Horticultural Crops: Environmental Implications and Thrusts (pp. 172-174). Association for Advancement of Pest Management in Horticultural Ecosystems, Bangalore.
- Gahukar R T 1995 Neem in Plant Protection. Agri-Horticultural Publishing House, Nagpur.
- Gahukar R T 1998 Commercial and industrial aspects of neem-based pesticides. Pestology 22 (10): 5-14.
- Gahukar R T 2003 Issues pertaining to the patentability of biotechnological subject matter in Indian agriculture. J. Intel. Property Rights 8: 9-22.
- Gopakumar B, Kumaresan D, Chandrasekhar S & Varadarasan S 1996 Evaluation of neem derivatives and *Lantana camara* L. against cardamom pests. In: Singh R P, Chari M S, Raheja A K & Krans W (Eds.) Neem & Environment (pp. 589-594). Oxford & IBH Publishing Company Ltd., New Delhi.
- Gupta R B L, Sharma S & Yadava C P S 2002 Compatibility of two entomofungi, *Metarrhizium anisopliae* and *Beauveria bassiana* with certain fungicides, insecticides and organic manures. Indian J. Entomol. 64: 48-52.
- Gupta V K, Ahlawat S P, Kumar R V & Datta A A 2010 Effect of season and year on azadirachtin A and oil content in neem (*Azadirachta indica* A. Juss.) seeds and relationship of azadirachtin A and oil content with rainfall, temperature and humidity. Curr. Sci. 99: 953-956.
- (IIHR) Indian Institute of Horticultural Research 2008 Annual Report for 2007-08. Indian Institute of Horticultural Research, Bangalore.
- (IISR) Indian Institute of Spices Research 2003 Annual Report for 2002-03, Indian Institute of Spices Research, Calicut.
- Jain K L & Ameta O P 2006 Bioefficacy of imidacloprid and betacyfluthrin against insect pests of chilli. Pestology 30 (1): 36-39.
- Karthikeyan M, Bhaskaran R, Radhika K, Mathiyazhagan S, Sandoskumar R, Alice D & Velazhahan B 2006 Biological control of leaf blight of onion by use of oil formulation. Indian J. Plant Prot. 34: 89-93.
- Keisa J T & Varatharajan R 1995 Efficacy of two neem products in the field control of *Scirtothrips dorsalis* Hood (Thysanoptera) on *Capsicum annum*. Indian J. Plant Prot. 23: 166-168.
- Kumar A K, Naik M K, Allolli T B & Hosmani R M 2005a Evaluation of genotypes, fungicides and plant extracts for the management of aflatoxin contamination in chilli caused by *Aspergillus flavus* Link. Fres. Indian J. Plant Prot. 33: 115-118.
- Kumar S, Mani C & Singh R N 2005b Field evaluation of omite and some other acaricides against broad mite, *Polyphagotarsonemus latus* (Banks.) on chilli under Varanasi condition of eastern Uttar Pradesh. Pestology 29 (11): 61-63.
- Lakshmisubramanian S, Nalini K, Muthuraman C, Sakthivel R, Sundara Balaji N & Murali P M 1998 Residue studies with azadirachtin formulations on bhendi, chilli and brinjal. Pestology 22 (8): 37-45.
- Lal R J 1993 Grow neem tree to sustain ecological balance. Kisan World 20 (4): 15-17.
- Leela N K & Ramana K V 2000 Nematicidal activity of the essential oil of allspice (*Pimenta dioica* L. Merr.). J. Plant Biol. 27: 75-76.
- Lalnuntluanga J & Singh H K 2008 Performance of certain chemicals and neem formulations against ginger shoot borer (*Dichocrocis punctiferalis* Guen.). Indian J. Entomol. 70 (2): 182-186.
- Mezhatsu R, Daiho L & Upadhyay D N 2008 Integrated management of rhizome rot of ginger caused by *Fusarium oxysporum* f. sp. *zingiber*. J. Ecofriendly Agric. 3 (1): 75-77.
- More S A, Kabre G B & Saindane Y S 2005 Effect of various insecticides on larval period of *Chrysoperla carnea* (Stephens). J. Insect Sci. 18: 111-113.
- Naik D J, Tyagaraj N E, Dinesh Kumar M, Madaiah D & Belavadi V V 2004 Effect of neem based insecticides for the control of thrips and shoot and capsule borer of cardamom. In: Abstracts of pa-

- pers, Symposium on Spices and Aromatic Crops: Commercialization of Spices, Medicinal and Aromatic Crops, 1-2 November 2004, Calicut (pp. 24-25), Indian Society for Spices, Calicut.
- Nakat R V, Khaire V M & Dethe M D 2000 Management of red vegetable mite (*Tetranychus neocaledonicus* Andre) infesting betelvine crop in India. *Pestology* 24 (6): 48-50.
- Pandey J C, Kumar R & Gupta R C 1992 Possibility of biological control of rhizome rot of ginger by different antagonists. *Prog. Hort.* 24: 227-232.
- Raj S T, Christopher D & Usha Rani S 2006 Management of fruit rot of chilli with different plant products. *Indian J. Plant Prot.* 34: 274-275.
- Ramarethinam S & Loganathan S 2001 Studies on the biology and management of swallow tail butterfly, *Papilio demoleus* L. (Lepidoptera: Papilionidae) infesting the curry leaf, *Murraya koenigii* (L.) Sprengel. *Pestology* 25 (12): 9-11.
- Ramarethinam S & Rajagopal B 1999 Efficacy of *Trichoderma* sp. organic amendments and seed dressing fungicides on rhizome rot of turmeric. *Pestology* 23 (4): 21-22.
- Rao M N, Muralidhara Rao G & Tirumala Rao D 1998 Efficacy of neem products and their combinations against chilli pod borers. *Andhra Agric. J.* 45: 179-181.
- Rao M N, Muralidhara Rao G & Tirumala Rao K 1999a Efficacy of neem products and their combinations against chilli thrips, *Scirtothrips dorsalis* Hood. *Pestology* 23 (3): 10-12.
- Rao M N, Muralidhara Rao G & Tirumala Rao K 1999b Efficacy of neem products and their combinations against *Aphis gossypii* Glover on chillies. *Andhra Agric. J.* 46: 122-123.
- Rao M S 2007 Biopesticides for management of nematodes in horticultural crops. *Indian J. Plant Prot.* 35: 202-205.
- Saha A K, Nath P S, De B K & Mohasin M D 2005 Effect of different insecticides on the control of whitefly (*Bemisia tabaci* Genn.) population, incidence of chilli leaf curl virus and yield of chilli in the plains of West Bengal. *Pestology* 29 (10): 24-27.
- Sekhar N C & Ram G M 2004 Study on the bioefficacy of oxymatrine, a plant derived pesticide vs. conventional pesticides in the management of pod borers of chilli (*Capsicum annum* L.). *Pestology* 28 (5): 8-10.
- Selvan M T, Gopalkumar B, Chandrasekharm S S, Singh J, Kumaresan D & Varadarasan S 1996 Survey on the occurrence of coccids infesting black pepper in higher altitudes of Idukki district, Kerala, and adaptive trial for their management. *Pestology* 20 (1): 19-23.
- Senguttuvan K, Kuttulam S, Manoharan T & Gunasekaran K 2005 Investigation of azadirachtin content in *Azadirachta indica* A. Juss ecotypes of Tamil Nadu. *Pestology* 29 (1): 34-36.
- Shanker J S & Parmar B S 1999 Recent developments in botanicals and biopesticides. *Indian J. Plant Prot.* 27: 139-154.
- Sharma B K 1998 Antifungal properties of biocontrol agents and plant extracts against causal fungi of yellows and rhizome rot of ginger. *J. Biol. Contr.* 12: 77-80.
- Simmonds M S J, Manlove J D, Blaney W M & Khambay B S 2000 Effect of botanical insecticides on the foraging and feeding behaviour of the coccinellid predator, *Cryptolaemus montrouzieri*. *Phytoparasitica* 28: 99-107.
- Sivaprakasam K & Seetharaman K 1997 Plant products for the management of plant diseases. *Kisan World* 24 (7): 49-50.
- Sivasithambaram K, Smith L D J & Goss O M 1981 Effect of potting media containing fresh saw dust and composted tree barks on *Phytophthora cinnamomi* Rands. *Austr. Plant Pathol.* 10: 20-21.
- Suresh D R, Karoshi V R & Hegde G V 2001 Indigenous knowledge for plant protection. *Indian Farmers Dig.* 34 (3): 31, 44.
- Venugopal M N 1999 Characterization, early detection and management of kokke kandu disease of cardamom. Final Report of Ad-hoc Research Scheme, Indian Institute of Spices Research, Calicut.
- Vir S 2007 Neem; Genetic diversity in India and its use as biopesticide and biofertilizer. *Indian J. Plant Prot.* 35: 185-193.
- Vir S & Yadav S S 2007 Bioefficacy of neem oil in the management of *Myzus persicae* (Sulz.): A serious pest of important spice crop (Cumin) in arid region. *Pestology* 31(1): 42-45.